



Risk of Stroke in National Registry of Atrial Fibrillation (NRAF) Participants, Stratified by CHADS₂ Score

CHADS₂ Score*	No. of Patients (n=1733)	No. of Strokes (n=94)	NRAF Crude Stroke Rate per 100 Patient-Years	NRAF Adjusted Stroke Rate, (95% CI)[†]
0	120	2	1.2	1.9 (1.2–3.0)
1	463	17	2.8	2.8 (2.0–3.8)
2	523	23	3.6	4.0 (3.1–5.1)
3	337	25	6.4	5.9 (4.6–7.3)
4	220	19	8.0	8.5 (6.3–11.1)
5	65	6	7.7	12.5 (8.2–17.5)
6	5	2	44.0	18.2 (10.5–27.4)

*CHADS₂ score is calculated by adding 1 point for each of the following conditions: recent congestive heart failure, hypertension, age at least 75 years, and diabetes mellitus; and adding 2 points for having had a prior stroke or transient ischemic attack.

[†]The adjusted stroke rate is the expected stroke rate per 100 patient-years from the exponential survival model, assuming that aspirin was not taken.

Reproduced with permission from: Gage BF, Waterman AD, Shannon W, Boehler M, Rich MW, Radford MJ. Validation of clinical classification schemes for predicting stroke: results from the National Registry of Atrial Fibrillation. *JAMA*. 2001;285:2867.

©2006. American College of Physicians. Medical Knowledge Self-Assessment Program (MKSAP®14)