

## Practical Gynecology, 2<sup>nd</sup> Edition – Chapter 1 References

1. **Frye CA, Weisberg RB.** Increasing the incidence of routine pelvic examinations: behavioral medicine's contribution. *Womens Health.* 1994;21:33-55.
2. **Rodriquez MA, Bauer HM, McLoughlin E, et al.** Screening and interventions for intimate partner abuse: practices and attitudes of primary care physicians. *JAMA.* 1999;282:468-74. *In a survey of California physicians, most had cared for patients who had experienced abuse.*
3. **Warshaw C, Alpert E.** Integrating routine inquiry about domestic violence into daily practice. *Ann Intern Med.* 1999;131:610-2.
4. **Saunders KJ, Pilgrim CA, Pennypacker HS.** Increased proficiency in breast self-examination. *Cancer.* 1986;58:2531-7.
5. **Ehrenthal DB, Farber NJ, Collier VU, Aboff BM.** Chaperone use by residents during pelvic, breast, testicular and rectal exams. *J Gen Intern Med.* 2000;15:573-6. *One of the few surveys of physicians on this topic.*
6. **Goswamy RK, Campbell S, Royston JP, et al.** Ovarian size in postmenopausal women. *Br J Obstet Gynaecol.* 1988;95:795-801.
7. **Goswamy RK, Campbell S, Whitehead MI.** Screening for ovarian cancer. *Clin Obstet Gynecol.* 1983;10:621.
8. **Griffith WF, Stuart GS, Gluck KL, Heartwell SF.** Vaginal speculum lubrication and its effects on cervical cytology and microbiology. *Contraception.* 2005;72:60-4.
9. **Harer WB, Valenzuela G Jr, Lebo D.** Lubrication of the vaginal introitus and speculum does not affect Papanicolaou smears. *Obstet Gynecol.* 2002;100(5 Pt 1):887-8.
10. **Amies AM, Miller L, Lee SK, Koutsky L.** The effect of vaginal speculum lubrication on the rate of unsatisfactory cervical cytology diagnosis. *Obstet Gynecol.* 2002;100(5 Pt 1):889-92.
11. **Pearce KF, Haefner HK, Sarwar SF, Nolan TE.** Cytopathological findings on vaginal Papanicolaou smears after hysterectomy for benign gynecologic disease. *N Engl J Med.* 1996;335:1559-62. *Well-done study that has changed medical practice.*
12. **American Geriatrics Society.** Screening for cervical carcinoma in older women. *J Am Ger Soc.* 2001;49:655-7.
13. **Maryrand MH, Duarte-Franco E, Rodrigues I, et al.** Human papillomavirus DNA versus Papanicolaou screening tests for cervical cancer. *N Engl J Med.* 2007;357:1579-88.
14. **Gaydos, CA, Howell MR, Pare B, et al.** Ligase chain reaction screening for chlamydia yielded a high prevalence of chlamydial infection in female military recruits.

N Engl J Med. 1998;229:739-44. *Younger age was a major risk factor for chlamydial infection.*

15. **Engstrum JL, Ostrenga KG, Plass RV, Work BA.** The effect of maternal bladder volume on fundal height measurements. Br J Obstet Gynaecol. 1989;96:987-91.

16. **Worthen N, Bustillo M.** Effect of urinary bladder fullness on fundal height measurements. Am J Obstet Gynecol. 1980;138:759-62.

17. **Nadel MR, Shapiro JA, Klabunde CN et al.** A national survey of primary care physicians' methods for screening for occult blood. Ann Intern Med. 2005;142:86-94. *It is most important that a positive stool guaiac receives further evaluation, especially while acceptance of colon cancer screening is suboptimal.*

18. **Koninckx PR, Meuleman C, Oosterlynck D, Cornillie FJ.** Diagnosis of deep endometriosis by clinical examination during menstruation and plasma CA-125 concentration. Fertil Steril. 1996;65:280-7. *The gold standard for diagnosis of endometriosis remains laparoscopy; however, because endometrial implants are most enlarged during menstruation, they may be more palpable at that time.*

19. **Koadlow E.** The pelvic examination: the first vaginal examination for a healthy young woman. Aust Fam Physician. 1990;19:665-9. *Thoughtful review.*

20. **Rimsza ME.** An illustrated guide to adolescent gynecology. Pediatr Clin North Am. 1989;36:639-63.

21. **Walsh JM, Dolan NC, Charney P, et al.** Update in women's health. Ann Intern Med. 2000;133:808-14.

22. **Mandblatt J, Gopaul I, Wistreich M.** Gynecological care of elderly women: another look at Papanicolaou smear testing. JAMA. 1986;256:367-93. *Not having received regular gynecologic care was a major risk factor for an abnormal Pap smear result.*

23. **Milsom I, Arvidsson L, Ekelund P, et al.** Factors influencing vaginal cytology, pH and bacterial flora in elderly women. Acta Obstet Gynecol Scand. 1993;72:286-91.

24. **Wee CC, McCartney EP, Davis RB, Phillips RS.** Screening for cervical and breast cancer: is obesity an unrecognized barrier to preventive care? Ann Intern Med. 2000;132:697-704. *Obese women were less likely to receive what are considered standard health maintenance examinations and tests.*