

Practical Gynecology, 2nd Edition – Chapter 9 References

*1. **Goldner TE, Lawson HW, Xia Z, Atrash HK.** Surveillance for ectopic pregnancy--United States, 1970-1989. *MMWR CDC Surveill Summ.* 1993;42:73-85.

*2. **Seeber BE, Barnhart KT.** Suspected ectopic pregnancy. *Obstet Gynecol.* 2006;107(2 Pt 1):399-413. *A new hCG “doubling” time was recently established, and this study reviews the diagnosis and management of ectopic pregnancy using this new parameter. The minimal rise in hCG in a viable pregnancy is 53% in 2 days, while the minimal decline of a spontaneous abortion is 21% to 35% over 2 days (depending on the initial level). A rise or fall in serial hCG values that is slower than this is suggestive of an ectopic pregnancy.*

*3. **Tay JI, Moore J, Walker JJ.** Ectopic pregnancy. *BMJ.* 2000;320:916-9.

*4. **Saraiya M, Berg CJ, Kendrick JS, Strauss LT, Atrash HK, Ahn YW.** Cigarette smoking as a risk factor for ectopic pregnancy. *Am J Obstet Gynecol.* 1998;178:493-8.

*5. **Stovall TG, Kellerman AL, Ling FW, Buster JE.** Emergency department diagnosis of ectopic pregnancy. *Ann Emerg Med.* 1990;19:1098-103.

*6. **Atri M, Bret PM, Tulandi T.** Spontaneous resolution of ectopic pregnancy: Initial appearance and evolution at transvaginal US. *Radiology.* 1993;186:83-6.

7. **Ushakov FB, Elchalal U, Aceman PJ, Schenker JG.** Cervical pregnancy: Past and future. *Obstet Gynecol Surv.* 1997;52:45-59.

*8. **Karande VC, Flood JT, Heard N, Veeck L, Muasher SJ.** Analysis of ectopic pregnancies resulting from in-vitro fertilization and embryo transfer. *Hum Reprod.* 1991;6:446-9.

*9. **Weckstein LN, Boucher AR, Tucker H, Gibson D, Rettenmaier MA.** Accurate diagnosis of early ectopic pregnancy. *Obstet Gynecol.* 1985;65:393-7.

*10. **Seeber BE, Sammel MD, Guo W, Zhou L, Hummel A, Barnhart KT.** Application of redefined human chorionic gonadotropin curves for the diagnosis of women at risk for ectopic pregnancy. *Fertil Steril.* 2006;86:454-9.

*11. **American College of Obstetricians and Gynecologists.** Medical management of tubal pregnancy. *ACOG Practice Bulletin.* 1998;3:4. *Available at www.acog.org.*

11.A **Seeber BE, Barnhart KT.** Suspected ectopic pregnancy. *Obstet Gynecol.* 2006;107(2 Pt 1):399-413.

*12. **Kadar N, Bohrer M, Kemmann E, Shelden R.** The discriminatory human chorionic gonadotropin zone for endovaginal sonography: A prospective, randomized study. *Fertil Steril.* 1994;61:1016-20.

- *13. **Morlock RJ, Lafata JE, Eisenstein D.** Cost-effectiveness of single-dose methotrexate compared with laparoscopic treatment of ectopic pregnancy. *Obstet Gynecol.* 2000;95:407-12.
14. **Sherman D, Langer R, Sadovsky G, Bukovsky I, Caspi E.** Improved fertility following ectopic pregnancy. *Fertil Steril.* 1982;37:497-502.
- *15. **Shannon C, Brothers LP, Philip NM, Winikoff B.** Infection after medical abortion: A review of the literature. *Contraception.* 2004;70:183-90.
- *16. **Cohen AL, Bhatnagar J, Reagan S, Zane SB, D'Angeli MA, Fischer M, et al.** Toxic shock associated with *Clostridium sordelli* and *Clostridium perfringens* after medical and spontaneous abortion. *Obstet Gynecol.* 2007;110:1027-33. *This CDC study analyzed four cases of toxic shock caused by Clostridium associated with medical or spontaneous abortion, and determined that two were caused by C. perfringens and two by C sordelii. These patients did not have fever, but presented with weakness plus nausea/vomiting or diarrhea (with or without abdominal pain). Women who have undergone medical abortion or have recently been pregnant presenting with such symptoms should be evaluated for sepsis.*
- *17. **American College of Obstetricians and Gynecologists.** Medical management of abortion. *ACOG Practice Bulletin.* 2005;67:3.
- *18. **Haggerty CL, Ness RB.** Newest approaches to treatment of pelvic inflammatory disease: A review of recent randomized clinical trials. *Clin Infect Dis.* 2007;44:953-60.
- *19. **McNeeley SG, Hendrix SL, Mazzoni MM, Kmak DC, Ransom SB.** Medically sound, cost-effective treatment for pelvic inflammatory disease and tubo-ovarian abscess. *Am J Obstet Gynecol.* 1998;178:1272-8.
- *20. **Centers for Disease Control and Prevention.** Updated recommended treatment regimens for gonococcal infections and associated conditions--United States, April 2007. Division of STD Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD and TB Prevention. 5 Jun 2007. Available from: www.cdc.gov/std/treatment/2006/updated-regimens.htm. *In 2007, the CDC released updated treatment guidelines for gonorrhea due to widespread fluoroquinolone resistance, and recommended the use of cephalosporins instead. For patients with severe cephalosporin allergy and uncomplicated gonorrhea, spectinomycin can be substituted. For other situations, infectious diseases consultation may be helpful.*
- *21. **Centers for Disease Control and Prevention.** Sexually transmitted diseases treatment guidelines, 2006. *MMWR.* 2006;55:56-61, 80-86.
- *22. **Ness RB, Soper DE, Holley RL, Peipert J, Randall H, Sweet RL, et al.** Effectiveness of inpatient and outpatient treatment strategies for women with pelvic inflammatory disease: Results from the pelvic inflammatory disease evaluation and clinical health (PEACH) randomized trial. *Am J Obstet Gynecol.* 2002;186:929-37. *The*

PEACH trial was a landmark RCT that demonstrated that outcomes of PID of mild-to-moderate severity were similar whether patients were treated as inpatients or outpatients. Longer follow-up (mean of 35 months) compared parameters such as pregnancy rates, time to pregnancy, PID recurrence, chronic pelvic pain or ectopic pregnancy, and found them to be equal in the two groups. Since most women with PID are managed as outpatients, this study is very reassuring.

23. **Aziz D, Davis V, Allen L, Langer JC.** Ovarian torsion in children: Is oophorectomy necessary? *J Pediatr Surg.* 2004;39:750-3.

*24. **Houry D, Abbott JT.** Ovarian torsion: A fifteen-year review. *Ann Emerg Med.* 2001;38:156-9. *This retrospective chart review examined 87 patients diagnosed with ovarian torsion. Most described the pain as “sharp” or stabbing and had nausea and vomiting, but otherwise the presentations varied, and only 3 had peritoneal signs. Not surprisingly, less than one third of patients underwent surgery within 24 hours, and detorsion was possible in only 9% of patients.*

25. **Servaes S, Zurakowski D, Laufer MR, Feins N, Chow JS.** Sonographic findings of ovarian torsion in children. *Pediatr Radiol.* 2007;37:446-51.

26. **Varras M, Tsikini A, Polyzos D, Samara C, Hadjopoulos G, Akrivis C.** Uterine adnexal torsion: Pathologic and gray-scale ultrasonographic findings. *Clin Exp Obstet Gynecol.* 2004;31:34-8.

27. **Albayram F, Hamper UM.** Ovarian and adnexal torsion: Spectrum of sonographic findings with pathologic correlation. *J Ultrasound Med.* 2001;20:1083-9.

*28. **Chen KT, Huard RC, Della-Latta P, Saiman L.** Prevalence of methicillin-sensitive and methicillin-resistant *Staphylococcus aureus* in pregnant women. *Obstet Gynecol.* 2006;108(3 Pt 1):482-7.

*29. **Jamart S, Denis O, Deplano A, Tragas G, Vanderghenst A, De Bels D, et al.** Methicillin-resistant staphylococcus aureus toxic shock syndrome. *Emerg Infect Dis.* 2005;11:636-7.

*30. **Hajjeh RA, Reingold A, Weil A, Shutt K, Schuchat A, Perkins BA.** Toxic shock syndrome in the united states: Surveillance update, 1979 1996. *Emerg Infect Dis.* 1999 Nov-Dec;5(6):807-10.

*31. **Centers for Disease Control and Prevention.** Toxic shock syndrome. Centers for Disease Control and Prevention. 24 Oct 2005. Available from: www.cdc.gov/ncidod/dbmd/diseaseinfo/toxicshock_t.htm.

32. **American Medical Association.** Strategies for treatment and prevention of sexual assault. Chicago: American Medical Association; 1995.

*33. **American College of Obstetricians and Gynecologists.** Special issues: Sexual assault. In: Special Issues in Women's Health. Washington, DC: American College of Obstetricians and Gynecologists; 2005:189-199.

34. **Marchbanks PA, Lui KJ, Mercy JA.** Risk of injury from resisting rape. Am J Epidemiol. 1990 Sep;132(3):540-9.

35. **Centers for Disease Control and Prevention.** Case-control study of HIV seroconversion in health care workers after percutaneous exposure to HIV-infected blood: France, United Kingdom, and United States, January 1988-August 1994. MMWR. 1995;44:929-933.

36. **Burgess AW, Holmstrom LL.** Rape trauma syndrome. In: Rape: Victims of Crisis. Bowie, MD: Robert J Brady Co; 1974:37-50.