

## **Practical Gynecology, 2<sup>nd</sup> Edition – Chapter 10 References**

- \*1. **Thom D.** Variation in estimates of urinary incontinence prevalence in the community: effects of differences in definition, population characteristics, and study type. *J Am Geriatr Soc.* 1998;46:473-80.
2. **Herzog AR, Fultz NH.** Prevalence and incidence of urinary incontinence in community-dwelling populations. *J Am Geriatr Soc.* 1990;38:273-81.
3. **Hohide E, Pringle D, Robertson D, Chambers L.** Prevalence of urinary incontinence in patients receiving home services under Medicare. *Health Care Finance Rev.* 1988;10:105-8.
4. **Ouslander JG, Zarit SH, Orr NK, Muira SA.** Incontinence among elderly community-dwelling dementia patients: characteristics, management and impact on caregivers. *J Am Geriatr Soc.* 1990;38:440-5.
5. **Diokno AC, Brock BM, Brown, MB, et al.** Prevalence of urinary incontinence and other urologic symptoms in the non-institutionalized elderly. *J. Urol.* 1986;136:1022.
6. **Johnson MJ, Werner C.** We have no choice: a study of familial guilt feelings surrounding nursing home care. *J Geront Nurs.* 1982;8:641-5,654.
7. **Ouslander JG, Kane RL, Abrass IB.** Urinary incontinence in elderly nursing home patients. *JAMA.* 1982;248:1194.
8. **Starer P, Libow LS.** Obscuring urinary incontinence. *J Am Geriatr Soc.* 1985;33:842.
- \*9. **Wolin LH.** Stress incontinence in young, healthy nulliparous females. *J Urol.* 1969;101:545.
10. **Cardozo L.** Urinary incontinence in women: have we anything new to offer? *BMJ.* 1991;303:1453.
11. **Norton P, MacDonald L, Stanton S.** Distress associated with female urinary complaints and delay in seeking treatment. *Neurourol Urodyn.* 1987;6:170.
- 11A. **Hendrix SL, Cochrane BB, Nygaard IE, et al.** Effects of estrogen with and without progestin on urinary incontinence. *JAMA.* 2005;293:935-48.
12. **Wyman JF, Harkins SW, Fantl JA.** Psychosocial impact of urinary incontinence in the community-dwelling population. *J Am Geriatr Soc.* 1990;38:282-8.
13. **Dugan E, Cohen SJ, Bland DR, et al.** The association of depressive symptoms and urinary incontinence among older adults. *J Am Geriatr Soc.* 2000;48:413-6.

14. **Johnson TM II, Kincade JE, Bernard SL, et al.** The association of urinary incontinence with poor self rated health. *J Am Geriatr Soc.* 1998;46:693-9.
- \*15. **Temml C, Haidinger G, Schmidbauer J, et al.** Urinary incontinence in both sexes: prevalence rates and impact on quality of life and sexual life. *Neurourol Urodyn.* 2000;19:259-71.
- \*16. **Brown JS, Vittinghoff E, Wyman JF, et al.** Urinary incontinence: does it increase risk for falls and fractures? *J Am Geriatr Soc.* 2000;48:721-5.
- \*17. **Milsom I, Abrams P, Cardozo L, et al.** How widespread are the symptoms of an overactive bladder and how are they managed? A population-based prevalence study. *British J of Urol.* 2001;87:760-6.
18. **Nelson R, Furner S, Jesudason V.** Urinary incontinence in Wisconsin skilled nursing facilities: prevalence and associations in common with fecal incontinence. *J Aging Health.* 2001;13:539-47.
19. **Maggi S, Minicuci N, Langlois J, et al.** Prevalence rate of urinary incontinence in community-dwelling elderly individuals: the Veneto study. *J Gerontol A Biol Sci Med Sci.* 2001;56:M14-18.
- \*20. **Finkelstein MM.** Medical conditions, medications, and urinary incontinence: analysis of a population-based survey. *Can Fam Physician.* 2002;48:96-101.
21. **Fultz NH, Herzog AR, Raghunathan TE, et al.** Prevalence and severity of urinary incontinence in older African American and Caucasian women. *J Gerontol A Biol Sci Med Sci.* 1999;54:M299-M303.
22. **Abrams P, Cardozo L, Fall M, et al.** The standardisation of terminology in lower urinary tract function: report from the standardization sub-committee of the International Continence Society. *Urology.* 2003;61:37-49.
23. **Stamey TA.** Urinary incontinence in the female. In: Walsh PC, Retik AB, Stamey TA, Vaughan DE Jr., eds. *Campbell's Urology*, 6th ed. Philadelphia; WB Saunders; 1992:2829-50.
24. **McGuire EJ, Lytton B, Pepe V, et al.** Stress incontinence. *Am J Obstet Gynec.* 1976;47:255.
25. **McGuire EJ, Savastano JA.** Stress incontinence and detrusor instability/urge incontinence. *Neurourol Urodyn.* 1985;4:313.
26. **Farrar DJ, Osborne JL, Stephenson TP, et al.** A urodynamic view of bladder outflow obstruction in the female: factors influencing the results of treatment. *Br J Urol.* 1976;47:815.
27. **Axelrod SL, Blaivas JG.** Bladder neck obstruction in women. *J Urol.* 1987;137:497.

28. **Stephens FD.** Anatomic vagaries of double ureters. *Aust N Z J Surg.* 1958;28:27.
29. **Ellerker AG.** The extravesical ectopic ureter. *Br J Surg.* 1958;45:344.
30. **Hinman F Jr, Cohlman WR.** Gartner's duct carcinoma in a urethral diverticulum. *J Urol.* 1960;83:414.
31. **Benjamin J, Eliot L, Cooper JF, et al.** Urethral diverticulum in adult females: clinical aspects, operative procedures, and pathology. *Urology.* 1974;3:1.
32. **Peters WH, Vaughan ED Jr.** Urethral diverticulum in the female. *Obstet Gynecol.* 1976;47:549.
33. **Leach GE, Bavendam TG.** Female urethral diverticula. *Urology.* 1987;30:110.
- \*34. **Hilton P.** Urinary incontinence during sexual intercourse: a common, but rarely volunteered symptom. *Br J Obstet Gynecol.* 1988;95:377.
- \*35. **Khan Z, Bhola A, Starer P.** Urinary incontinence during orgasm. *Urology.* 1988;31:279.
- \*36. **Snooks SJ, Swash M, Henry MM, et al.** Risk factors in childbirth causing damage to pelvic floor innervations. *Int J Colorect Dis.* 1986;1:20.
37. **Smith ARB, Hosker GL, Warrell DW.** The role of partial denervation of the pelvic floor in the aetiology of genitourinary prolapse and stress incontinence of urine: a neurophysiological study. *Br J Obstet Gynecol.* 1989;96:24.
- \*37A. **Altman D, Granath F, Cnattingius S, Falconer C.** Hysterectomy and risk of stress-urinary-incontinence surgery: nationwide cohort study. *Lancet.* 2007;370:1494-9.
- \*37B. **Kuruba R, Almahmeed T, Martinez F, Torrella TA, Haines K, Nelson LG, Gallagher SF, Murr MM.** Bariatric surgery improves urinary incontinence in morbidly obese individuals. *Surg Obes Relat Dis.* 2007 Nov-Dec;3(6):586-90; discussion 590-1.
38. **Blaivas JG, Zayed AAH, Labib KB.** The bulbocavernous reflex in urology: a prospective study of 299 patients. *J Urol.* 1981;126:197.
- \*39. **Schmidt RA, Senn E, Tanagho EA.** Functional evaluation of sacral nerve root integrity. *Urology.* 1990;35:5.
40. **Marshall FC, Uson AC, Melicow MM.** Neoplasms and caruncles of the female urethra. *Surg Gynecol Obstet.* 1960;110:723.
- \*41. **Fantl JA, Newman DK, Colling J, et al.** Urinary incontinence in adults: acute and chronic management. Rockville, MD: US Department of Health and Human Services, Agency for Health Care Policy and Research, March 1996; AHCPR Publication 96-0682, Clinical Practice Guideline No. 2, 1996 Update.

\*42. **Burgio KL, Locher JL, Goode PS, et al.** Behavioral vs. drug treatment for urge urinary incontinence in older women. *JAMA*. 1998;280:1995-2000. *This randomized placebo-controlled trial compared behavioral and drug therapy for urge and mixed incontinence, and found that behavioral treatment was more effective in reducing episodes of incontinence (mean 80.7% reduction compared to mean 68.5% reduction; P=0.04). Patient-perceived improvement was also greater for behavioral treatment, and only 14.0% of patients randomized to this group wished to change to another treatment.*

\*43. **Klarov P, Belving D, Bischiff N, et al.** Pelvic floor exercises versus surgery for female urinary stress incontinence. *Urol Int*. 1986;41:129.

44. **Mauritsen L.** Pelvic floor exercises for female stress urinary incontinence. *Int Urogynecol J*. 1994;5:44.

\*45. **Hay-Smith EJ, Dumoulin C.** Pelvic floor muscle training versus no treatment, or inactive control treatments, for urinary incontinence in women. *Cochrane Database of Systematic Reviews* (1). CD005654, 2006. *Pelvic floor muscle training (PFMT) is a commonly used physical therapy treatment for urinary incontinence. Overall, this review of data from 61 trials provides some support for the widespread recommendation that PFMT be included as first-line conservative management for women with stress, urge, or mixed urinary incontinence. Younger women (age 40s and 50s) with stress urinary incontinence alone, who participate in a supervised PFMT program for at least three months, may benefit the most.*

\*46. **Kegel A.** Progressive resisting exercises in the functional restoration of the perineal muscles. *Am J Obstet Gynecol*. 1949;56:238.

47. **Bo K, Talseth T, Home I.** Single blind, randomized controlled trial of pelvic floor exercises, electrical stimulation, vaginal cones, and no treatment in management of genuine stress incontinence in women. *BMJ*. 1999;318:487-93.

48. **Bo K, Llarger S, David S, et al.** Knowledge about and ability to correct pelvic floor muscle exercises in women with urinary stress incontinence. *Neurourol Urodyn*. 1988;71:261.

49. **Cardoza LD, Abrams PD, Stanton SL, et al.** Idiopathic bladder instability treated by biofeedback. *Br J Urol*. 1978;50:521.

50. **Stroebel CF, Glueck BC.** Biofeedback treatment in medicine and psychiatry; an ultimate placebo? *Sem Psychiatr*. 1973;5:379.

51. **Plevnick S.** New method for testing and strengthening of pelvic floor muscles. In: *Proceedings of the 15<sup>th</sup> Annual General Meeting, International Continence Society, 1985.*

52. **Abrams PH, Torrens M.** Urine flow studies. *Symposium on Clinical Urodynamics*. *Urol Clin North Am*. 1979;6:63.

53. **Farrar DJ, Whiteside CG, Osborne JL, et al.** A urodynamic analysis of micturition symptoms in the female. *Surg Gyn Obstet*. 1975;141:875.

\*54. **Nabi G, Cody JD, Ellis G, Herbison P, Hay-Smith J.** Anticholinergic drugs versus placebo for overactive bladder syndrome in adults. *Cochrane Database of Systematic Reviews* (4). CD003781, 2006. *Anticholinergic drugs are commonly used to treat overactive bladder (urge incontinence and/or (urgency with frequency). Overall, evidence from this review of 61 trials suggests the use of anticholinergic drugs results in statistically significant improvement in symptoms (decreased number of voids and leakage episodes) with modest improvement in quality of life. Dry mouth, a common side effect of therapy, did not lead to drug discontinuation. It is unknown whether benefits are sustained during long-term treatment or after treatment stops.*

55. **Khullar V, Cardozo LD, Abbott D, Anders K.** GAX collagen in the treatment of urinary incontinence in elderly women: a two year follow up. *Br J Obstet Gynaecol.* 1997;104:96-9.

\*56. **Culligan PJ, Heit M.** Urinary incontinence in women: evaluation and management. *Am Fam Physician.* 2000;62:2433-52. *This article provides the primary care physician with an overview of the basic evaluation and management of urinary incontinence.*

\*57. **Keegan PE, Atiemo K, Cody J, McClinton S, Pickard R.** Periurethral injection therapy for urinary incontinence in women. *Cochrane Database of Systematic Reviews* (3). CD003881, 2007. *This review provides limited data due to the small size and moderate quality of the 12 trials; meta-analysis was not possible. Injection therapy may be a useful option for short-term symptomatic relief in women with co-morbidity that precludes anesthesia. To achieve a satisfactory result two or three injections are likely to be required. Further research is needed.*

58. **Burch JC.** Urethrovaginal fixation to Cooper's ligament for correction of stress incontinence, cystocele, and prolapsed. *Am J Obstet Gynecol.* 1961;81:281.

\*59. **Albo ME, Richter HE, Brubaker L, Norton P, Kraus SR et al.** for the Urinary Incontinence Treatment Network. Burch Colposuspension versus Fascial Sling to Reduce Urinary Stress Incontinence. *N Engl J Med.* 2007;356:2143-55

\*60. **Gittes RF, Loughlin KR.** No-incision pubovaginal suspension for stress incontinence. *J Urol.* 1987;138:568.

\*61. **Silva WA.** Treatment of stress urinary incontinence--midurethral slings: top-down, bottom-up, "outside-in," or "inside-out". *Clin Obstet Gynecol.* 2007;50:362-75.