

Practical Gynecology, 2nd Edition – Chapter 21 References

- *1. **Saltzman LE, Fanslow JL, McMahon PM, Shelley GA.** Intimate Partner Violence Surveillance: Uniform Definitions and Recommended Data Elements. Atlanta: National Center for Injury Prevention and Control, Centers for Disease Control and Prevention; 1999. NCJ. 181867.
2. **Hathaway JE, Mucci LA, Silverman JG, et al.** Health status and health care use of Massachusetts women reporting partner abuse. *Am J Prev Med.* 2000;19:302-7.
- *3. **Wisner CL, Gilmer TP, Saltzman LE, Zink TM.** Intimate partner violence against women: do victims cost health plans more? *J Fam Pract.* 1999;48:439-443.
- *4. **Ulrich YC, Cain KC, Sugg NK, et al.** Medical care utilization patterns in women with diagnosed domestic violence. *Am J Prev Med.* 2003;24:9-15.
- *5. **Nicolaidis C.** The Voices of survivors documentary: using patient narrative to educate physicians about domestic violence. *J Gen Intern Med.* 2002;17:117-24.
6. **Johnson MP, Ferraro KJ.** Research on domestic violence in the 1990s: Making distinctions. *Journal of Marriage & the Family.* 2000;62:948-963.
- *7. **Nicolaidis C, Curry M, McFarland B, Gerrity M.** Violence, mental health and physical symptoms in an academic internal medicine practice. *J Gen Intern Med.* 2004;19:819-27.
8. **Tollestrup K, Sklar D, Frost FJ, et al.** Health indicators and intimate partner violence among women who are members of a managed care organization. *Prev Med.* 1999;29:431-440.
9. **Bauer HM, Rodriguez MA, Perez-Stable EJ.** Prevalence and determinants of intimate partner abuse among public hospital primary care patients. *J Gen Intern Med.* 2000;15:811-817.
- *10. **Coker AL, Smith PH, McKeown RE, King MJ.** Frequency and correlates of intimate partner violence by type: physical, sexual, and psychological battering. *Am J Public Health.* 2000;90:553-559.
11. **Tjaden P, Thoennes N, Allison CJ.** Comparing violence over the life span in samples of same-sex and opposite-sex cohabitants. *Violence Vict.* 1999;14:413-425.
- *12. **Tjaden P, Thoennes N.** Extent, nature, and consequences of intimate partner violence: findings from the National Violence Against Women Survey. Washington (DC): Department of Justice (US) 2000. Publication No. NCJ181867.
13. **Fox JA, Zawitz MW.** Homicide trends in the US. Washington, DC: Department of Justice (US). 2004.

14. **Wadman MC, Muelleman RL.** Domestic violence homicides: ED use before victimization. *Am J Emerg Med.* 1999;17:689-91.
- *15. **Kothari CL, Rhodes KV.** Missed opportunities: emergency department visits y police-identified victims of intimate partner violence. *Ann Emerg Med.* 2006; 47:190-9.
16. **McCauley J, Kern DE, Kolodner K, et al.** The “battering syndrome”: prevalence and clinical characteristics of domestic violence in primary care internal medicine practices. *Ann Intern Med.* 1995;123:737-746. *Of approximately 2000 female patients attending primary care community clinics, 5 % had experienced domestic violence in the prior year and 21% had experienced domestic violence at some time in her adult life. Recently abused women were more likely to visit the hospital and have more mental and physical symptoms and signs.*
- *17. **Campbell J, Jones AS, Dienemann J, et al.** Intimate partner violence and physical health consequences. *Arch Intern Med.* 2002;162(10):1157-63.
18. **Coker AL, Smith PH, Bethea L, et al.** Physical health consequences of physical and psychological intimate partner violence. *Arch Fam Med.* 2000;9:451-457. *The investigators found that compared to never abused women in primary care practice, women who had experienced either psychological or physical intimate partner violence had greater relative risks of multiple adverse health consequences including chronic pain (pelvic, abdominal), sexually transmitted infections, constipation or diarrhea, and headaches.*
- *19. **American Medical Association.** Report 7 of the Council of Scientific Affairs. Diagnosis and Management of Family Violence, June 2005. <http://www.ama-assn.org/ama/pub/category/15248.html>. Accessed on November 1, 2007.
20. **American College of Obstetrics and Gynecology.** Screening – Domestic Violence. http://www.acog.org/departments/dept_notice.cfm?recno=17&bulletin=585. Accessed November 1, 2007.
- *21. **U.S. Preventive Services Task Force.** Screening for family and intimate partner violence: recommendation statement [see comment]. *Ann Intern Med.* 2004;140:382-386.
22. **Nelson HD, Nygren P, McInerney Y, Klein J, US Preventive Services Task Force.** Screening women and elderly adults for family and intimate partner violence: a review of the evidence for the US Preventive Services Task Force. *Ann Intern Med.* 2004;140:387-396.
23. **Calonge N, Allan JD.** Screening for family violence: what’s an evidence-based doctor to do? Letter to the Editor. *Ann Intern Med.* 2004; 141:82-83.
24. **Roberts GL, Lawrence JM, Williams GM, Raphael B.** The impact of domestic violence on women’s mental health. *Aust N Z J Public Health.* 1998;22:796-801.

- *25. **Petersen R, Gazmararian J, Andersen Clark K.** Partner violence: implications for health and community settings. *Womens Health Issues.* 2001;11:116-125.
26. **Golding JM.** Intimate partner violence as a risk factor for mental disorders: a meta-analysis. *Journal of Family Violence.* 1999;14:99-132.
27. **Thompson MP, Kaslow NJ, Kingree JB, et al.** Partner abuse and posttraumatic stress disorder as risk factors for suicide attempts in a sample of low-income, inner-city women. *J Trauma Stress.* 1999;12:59-72.
28. **Brokaw J, Fullerton-Gleason L, Olson L, et al.** Health status and intimate partner violence: a cross-sectional study. *Ann Emerg Med.* 2002;39:31-38.
29. **Feder GS, Hutson M, Ramsay J, Taket AR.** Women exposed to intimate partner violence expectations and experiences when they encounter health care professionals: A meta-analysis of qualitative studies. *Arch Intern Med.* 2006;166:22-37. *This meta-analysis of 25 studies found several key constructs regarding the expectations and experiences of women exposed to intimate partner violence when they encountered health care professionals. While there was some variation of responses based on timing of interaction, relationship of patient to professional, and patient readiness, women consistently wanted health professionals to be knowledgeable, culturally sensitive, nonjudgmental, empathic and to ensure the patient's autonomy and confidentiality.*
30. **Rodriguez MA, Sheldon WR, Bauer HM, Perez-Stable EJ.** The factors associated with disclosure of intimate partner abuse to clinicians. *Journal of Family Practice.* 2001;50:338-344. *The investigators randomly selected 375 female patients with a history of intimate partner violence and found that direct questioning by health professionals was the best predictor of disclosure of intimate partner violence.*
31. **Nicolaidis C, XXXX.** In preparation/submission. **AUTHOR: PLEASE UPDATE THIS REFERENCE.**
32. **Rhodes KV, Frankel RM, Levinthal N, et al.** "You're not a victim of domestic violence, are you?" Provider patient communication about domestic violence. *Ann Intern Med.* 2007 Nov 6;147:620-7.
- *33. **Paranjape A, Liebschutz J.** STaT: a three-question screen for intimate partner violence. *J Womens Health.* 2003;12:233-239.
- *34. **Sherin KM, Sinacore JM, Li XQ, Zitter RE, Shakil A.** HITS: a short domestic violence screening tool for use in a family practice setting. *Fam Med.* 1998; 30:508-12.
- *35. **Brown JB, Lent B, Brett PJ, Sas G, Pederson LL.** Development of the Woman Abuse Screening Tool for use in family practice. *Fam Med.* 1996;28:422-8.
- *36. **McFarlane J, Parker B, Soeken K, Bullock.** Assessing for abuse during pregnancy. Severity and frequency of injuries and associated entry into prenatal care. *JAMA.* 1992;3176-8.

- *37. **Kyriacou DN, Anglin D, Taliaferro E, et al.** Risk factors for injury to women from domestic violence against women [see comments]. *N Engl J Med.* 1999;341:1892-1898.
- *38. **Nicolaidis C, Curry MA, Urlich Y, et al.** Could we have known? A qualitative analysis of data from women who survived an attempted homicide by an intimate partner. *J Gen Intern Med.* 2003;18:788-794.34.
39. **Campbell JC, Webster D, Koziol-McLain J, et al.** Risk factors for femicide in abusive relationships: results from a multi-site case control study. *Am J Public Health.* 2003;93:1089-1097. *In this case-control study, proxies, who were knowledgeable about the deceased victim of intimate partner homicide, were interviewed to elucidate risk factors for intimate partner homicide (femicide). The investigators found these risk factors: prior physical abuse (especially during pregnancy), gun availability, perpetrator unemployment, stalking, and prior threats of homicide or suicide by the perpetrator.*
40. **Weisz A, Tolman R, Saunders DG.** Assessing the risk of severe domestic violence: The importance of survivor's predictions. *J Interpers Violence.* 2000;15:75-90.
41. **Campbell JC, Koziol-McLain J, Webster D, et al.** Research results from a national study of intimate partner femicide: the danger assessment instrument. *NIJ Briefs.* 2002.
42. **Scholle SH, Rost KM, Golding JM.** Physical abuse among depressed women. *J Gen Intern Med.* 1998;13(9):607-13.
43. **Nicolaidis C, Curry M, Gerrity M.** Health care workers' expectations and empathy toward patients in abusive relationships. *J Am Board Fam Pract.* 2005;18:159-65.
44. **Zink T, Elder N, Jacobson J.** How children affect the mother/victim's process in intimate partner violence. *Arch Pediatr Adolesc Med.* 2003;157:587-592.
- *45. **Holt VL, Kernie MA, Lumley T, Wolf M, Rivara FP.** Civil protection orders and risk of subsequent police-reported violence. *JAMA.* 2002;288:589-594.
46. **Warshaw C, Ganley A.** Improving the Health Care Response to Domestic Violence: A Resource Manual for Health Care Providers, 2nd edition. San Francisco, Calif: The Family Violence Prevention Fund in collaboration with the Pennsylvania Coalition Against Domestic Violence. Available at: <http://endabuse.org/programs/display.php3?DocID=238>. Accessed August 29, 2007.