

Table 4. Univariate Analysis of Association between Clinical and Laboratory Variables and Mortality

Variable	Survivors (n = 235)	Deceased (n = 32)	P Value
Clinical characteristics			
Age, y	37 (18–75)	53 (34–96)	<0.001*
Men, %	37	55	0.06†
Significant medical history, n (%)	5 (2)	10 (31)	<0.001†
Laboratory variables on presentation‡			
Neutrophil count, $\times 10^9$ cells/L	3.7 (0.7–16.3)	5.4 (0.8–15.5)	<0.001*
Lymphocyte count, $\times 10^9$ cells/L	0.8 (0.2–1.8)	0.8 (0.3–1.8)	>0.2*
Activated partial thromboplastin time, s	32.9 (22–57)	33 (23–87)	0.033*
Creatinine level, $\mu\text{mol/L}$	67 (40–683)	84.5 (48–326)	<0.001*
Alanine aminotransferase level, U/L	27 (7–233)	35.5 (12–179)	0.07*
Lactate dehydrogenase level, $\mu\text{kat/L}$	3.7 (1.0–13.1)	4.4 (2.5–16.5)	0.004*
C-reactive protein level, mg/L	26.7 (6–256)	68.9 (16.2–305)	0.001*

* Mann–Whitney U test.

† Chi-square test.

‡ Values are expressed as the median (range).