

High Value Care

ConsumerReports[®]Health

ACP

AMERICAN COLLEGE OF PHYSICIANS
INTERNAL MEDICINE | *Doctors for Adults*[®]

Annals of Internal Medicine

Upper endoscopy for GERD

When you need it—and when you don't

People with heartburn often undergo a procedure called endoscopy to see if gastro-esophageal reflux disease (GERD) is the cause, and if it is, to check periodically for complications. But that's usually not necessary. Here's why.

It usually doesn't lead to better treatment.

Frequent heartburn could be caused by GERD, a condition in which acid in your stomach rises up into the esophagus. Left untreated, GERD can inflame the lining of the esophagus, causing a condition known as esophagitis. A few GERD patients develop Barrett's esophagus, a disorder that can, in rare cases, lead to cancer of the esophagus. Doctors can check for those problems with an endoscope, a lighted, flexible tube with a camera. But the procedure usually isn't necessary because the chance of esophageal cancer in heartburn sufferers is very low, affecting only about 1 in 2,500 patients older than 50 each year. Even among people with Barrett's, the risk of cancer is quite low. As a result, frequent endoscopies are usually unnecessary, and rarely change the treatments doctor offer.



It can pose risks.

Endoscopy is usually safe, but can cause bleeding or puncture the stomach or esophagus. The sedatives used can cause breathing difficulty, abnormal heart rhythms, and other problems. And incorrect findings can lead to needless tests and treatments. For example, people misdiagnosed with cancer might have part of the esophagus surgically removed, and those misdiagnosed with Barrett's might be denied insurance.

It can be expensive.

The procedure typically costs about \$1,130 plus a fee for anesthesia, according to HealthcareBlueBook.com. And even insured patients may have substantial co-pays and deductibles.

So when is it warranted?

People with GERD don't need endoscopy unless:

- Symptoms remain after four to eight weeks of a proton pump inhibitor (PPI) such as esomeprazole (Nexium), lansoprazole (Prevacid and generic), or omeprazole (Prilosec and generic).
- Severe inflammation of the esophagus persists despite two months of PPI treatment.
- Heartburn occurs with anemia, difficulty swallowing, weight loss, frequent vomiting, vomiting blood, or black or red bowel movements.
- You have frequent difficulty swallowing, plus a history of narrowing of the esophagus.

The test might also be appropriate if you are a man 50 or older with GERD symptoms for more than five years plus other risk factors such as smoking or obesity. Most people who have Barrett's need endoscopy once every three to five years. Those with early signs of cancer require it more often. And those without Barrett's usually don't need it repeated at all.

USING THIS INFORMATION

This information is provided for you to use in discussions with your health-care provider. The content is for educational use only and is not a substitute for professional medical advice, diagnosis, or treatment. Always seek the advice of a qualified health-care provider about any medical condition. Never disregard, avoid, or delay in obtaining medical advice from a health-care provider because of something you have read in this report. Use of this report is at your own risk. Consumer Reports, the American College of Physicians, Annals of Internal Medicine, and their distributors are not liable for any loss, injury, or other damage related to your use of this report. This report is intended solely for consumers' personal, noncommercial use and may not be altered or modified in any way or used in advertising, for promotion, or for any other commercial purpose. Special permission is granted to organizations participating in the Consumer Reports health communication program to disseminate free copies of this information in print or PDF formats to individual patients, members, or employees. Learn more at ConsumerHealthChoices.org or send an e-mail to HealthImpact@cr.consumer.org.

Published by Consumer Reports © 2012 Consumers Union of U.S., Inc., 101 Truman Avenue, Yonkers, NY 10703-1057. All rights reserved. Developed in cooperation with the American College of Physicians and Annals of Internal Medicine as part of the High Value Care patient education program. Portions of this report are derived from materials developed for the High Value, Cost-Conscious Care Initiative of the American College of Physicians and Annals of Internal Medicine and are used with permission. © 2012 American College of Physicians.


Consumer Reports' Advice

How should you manage heartburn?

The following steps can help relieve occasional heartburn as well as GERD:

• **Make lifestyle changes.**

Lose excess weight, stop smoking, eat smaller meals, don't lie down for several hours after eating, wear loose-fitting clothes, and limit how much alcohol you consume. Resist foods and beverages that can trigger your heartburn, such as chocolate, peppermint, coffee (with or without caffeine), garlic, onions, and fried, spicy, fatty, or tomato-rich food.



Sleep comfortably. Place wood blocks under your bedposts to raise the head of your bed 6 to 8 inches.

Control acid. Over-the-counter antacids such as Maalox, Mylanta, Rolaids, or Tums may ease heartburn. For a stronger acid-reducer, try an H₂ blocker such as famotidine (Pepcid and generic) or ranitidine (Zantac and generic).

Consider a PPI for GERD. Those drugs relieve symptoms and heal the lining of the esophagus in almost all sufferers. For best results, take them 30 to 60 minutes before a meal. Omeprazole and lansoprazole are sold over the counter, as low-cost generics. If symptoms don't improve after two to four weeks, ask your doctor about taking the dose twice a day. For our free Best Buy Drug report on PPIs, go to www.CRBestBuyDrugs.org and click on "heartburn."